

**Oral Dermatology: Common Findings
Approach to Diagnosis, Differential
Diagnosis and Treatment**

Ginat W. Mirowski, DMD, MD

Adjunct Associate Professor
Department of Oral Pathology Medicine
Radiology
Indiana University School of Dentistry

Volunteer Clinical Associate Professor
Department of Dermatology
Indiana University School of Medicine

Private practice

Conflict of Interest

- I have no conflicts of interest to report
- I may discuss off label use of medications
- The medications and treatment suggestions mentioned in this talk are based on my current use patterns or on the review of the literature
- Other treatment options may be available

2

Learning Objectives

Following this talk, the attendees will be able to

- Identify common oral findings, oral lesions, primary skin diseases as well as systemic disorders
- Discuss the differential diagnosis of red and white oral lesions and the approach to obtaining a diagnosis
- Describe therapeutic options for the management of erosive oral disorders

Complete Oral Exam Is A Subset Of The Mucocutaneous Exam*

-
-
- larynx, otic ie conjunctiva, esophagus,
- Vulva
- Vaginal → with speculum; wet mount
- Proper positioning and visualization
- Patient participation (use of mirror)

* Often requires consultation referral


Multidisciplinary Team

- | | |
|-----------------------|----------------------|
| • Nurse practitioner | • Urologist |
| • Physician assistant | • Psychologist |
| • Primary physician | • Sex therapist |
| • Gynecologist | • Physical therapist |
| • Dentist | • Marriage counselor |
| • Ophthalmologist | • Others |
| • Gastroenterologist | |

Routine Oral Examination Extraoral: Head & Neck

- Face, eyes & nose
- Lymph nodes
- Thyroid gland
- Major salivary glands
- Lips & angles of the mouth

Approach to the Patient; NOT



- Patient's head is not stable
- Physician is not in comfortable position
- Lighting may be too bright
- Tongue blade
- No gauze
- Not using gloves!!!!

Courtesy of WWW

The Physical Examination

- Recline patient at 45°
- Adequate lighting
 -
 -
 -
- Access
 -
-

Pitfalls of the Oral Examination

- Moisture alters the appearance of lesions
- Use a piece of gauze to:
 - dry off the mucosa
 - retract the tongue and lips
- Minor trauma alters the primary morphology
- Limited number of reaction patterns

Oral Cavity

- Mucosa is thin; Saliva is critical to health
- Oral → Necessary for ADL i.e. eating, drinking, speaking
 - Stimulate saliva
 - Dilutes concentration of topical medications
 - Decreases contact time
 - Much psychosocial implications; “baggage”

Lips

- Inspect
- Retract
- Palpate

Gingiva

- Scalloped
- Interdental Papillae - pointy

Buccal Mucosa

- Stensen's Duct (Parotid Gland)
 - saliva
 - Buccal bite line
 - Across from M²
- Horizontal bite line

Hard Palate and Maxillary Arch

Inspect and Palpate

- Incisive papilla
- Torus palatinus
- Rugae
- Minor salivary gland ducts

Oropharynx

- Uvula
- Palatoglossus fold
- Palatine tonsil
- Palatopharyngeal fold
- Waldeyer's Ring

Lingual Tonsil Waldeyer's Ring

- Mucosal lymphatic tissues (MALT) of nasopharynx and oropharynx
 - Palatine tonsils (lateral oropharynx)
 - Nasopharyngeal / Adenoids (posterior to)
 - Tubal tonsils (nasopharynx)
 - Lingual tonsils (dorsal tongue)

Tongue

-
- (mushroom-like) erythematous or pigmented
- Filiform papillae (hair-like) white or acquired exogenous pigment
- Circumvallate Papillae
 - site of oral cancer
 - Visual examination requires retraction of the tongue
 - Palpation - bimanual examination

Variants and Findings

Hooker's Lips Psychotria

Fordyce Spots (Condition)

- Minute yellow papules
- Incidence > with age
- Ectopic sebaceous glands
- Unassociated with hair follicles

Geographic Tongue

- Well circumscribed, circinate plaques
- Asymptomatic --> burning and itching
- Dorsal tongue> ventral tongue> other mucosal surfaces
- Raised white border (elongated filiform papillae)
- Loss of filiform & fungiform papillae

Geographic Tongue Benign Migratory Glossitis

- Unknown etiology
- Move daily or weekly
- Differential diagnosis: erosive lichen planus, coated tongue, burn, aphthous stomatitis
- Therapy: Supportive, topical corticosteroids

Mucocele
(Retention Cyst)

- Obstruction of salivary glands
- Asymptomatic swelling
- Translucent papule
- Myxoid exudate
- Recurrent
- Treatment --> marsupialization

Ranula

- Large, bluish mass
- Mucocele in floor of mouth
- May plunge through mylohyoid muscle into the neck
- May be life threatening !!!!!

Melkersson Rosenthal
(Granulomatous Cheilitis)

- -
 -
 -
 -
- small noncaseating granulomas

Stomatitis Nicotina
(smokers keratosis, smokers patches)

- On the hard palate
- Pipes/cigars> cigarettes
- Characteristic erythematous umbilicated papules
- Caused by tars and heat in tobacco smoke
- Hyperkeratosis / acanthosis of epithelium; inflammation of submucosa and minor salivary glands; squamous metaplasia of ducts
- Not premalignant but an opportunity to discuss risk of oropharyngeal CA

Clinical Challenges

White Plaques on Oral Mucosa
Differential Diagnosis

- Poor oral hygiene
- Morsicatio buccarum et labiorum
- Pseudomembranous candidosis
- Recurrent aphthous ulcers*
- Oral hairy leukoplakia
- Viral enanthem
- Lichen planus

* Often both red and white

Mononucleosis - Enanthem

- Epstein-Barr virus
- Prodrome of headache, malaise, fever and pharyngitis
- Incubation 33-49 days
- Exanthem occurs in 35% patients
- Accompanying Strep pharyngitis in 20-25%

Rimsza et al: Pediatr Clin North Am 2005

Morsicatio Buccarum et Labiorum (Chronic Nibbling of Cheeks and Lips)

- Physical reaction to chronic trauma
- Frequently symmetric
- Thickened shredded red and white areas
- May peel off
- Treatment: patient education, acrylic shield or OCD regimen

Aphthous Ulcers (RAU, RAS)

- Single or multiple
- Painful oval ulcers
- White gray fibrinous pseudomembrane
- Red border
- Prodrome +/-
- No blisters

Aphthous Ulcers (RAU, RAS)

Epidemiology

- Most common cause oral ulcers
- RAU affects 5-66% of the population in US
- 40% of selected groups of children have a history of RAU ulcers < 5 years of age
- 60% of professional students (dental and medical)
- ag
- Recurrences increase with immunosuppression or after smoking cessation

Recurrent Aphthous Ulcers

Clinical Presentation

- Three forms (minor, major and herpetiform)
- Minor (Mikulicz)
 - Most common (85%)
 - Single or multiple
 - Discrete round or oval
 - < 1 cm
 - Heal without scarring in 7-14 days

Aphthous Ulcers

Major (Sutton)

- Single >> multiple
- 10%
- Round or oval
- Soft palate, lips, oropharynx
- 2 -3 cm
- Heal with scarring
- Heal in 6 weeks to months
- More common in HIV

Aphthous Ulcers Herpetiform

- Rarest; resemble HSV
- <5%
- Grouped
- 1-2 mm
- Crops of 10-100
- Unkeratinized; rarely keratinized mucosa
- Heal in 7-14 days
- Do not scar

Conditions Associated With Oral Aphthae

- Stress
- Trauma: biting, dental appliances, brushing
- Xerostomia (40% have hypofunction - Dr. Zunt)
- Smoking cessation
- Endocrine factors (decrease progesterone in luteal phase; ulcers regress in pregnancy)
- Food allergies
- Sodium lauryl sulphate detergent emulsifier & surface cleanser)
- Drugs: Vioxx, Zolof, Remeron, Feverfew, xerostomic agents, alendronate and many others

Systemic Conditions Associated With Aphthous Ulcers 1/2

- Hematinic deficiency (up to 20%)
- Iron, vitamin B12, folate and rarely vitamin B-1, B-2, and B-6
- GI Malabsorption (3%)
- Celiac disease, dermatitis herpetiformis and gluten-sensitive enteropathy (HLA DRW10 and DQW1), Crohn disease, pernicious anemia

**Systemic Conditions
Associated With Aphthous Ulcers 2/2**

- Systemic lupus erythematosus
- HIV
- Behçet
- Trisomy 8, myelodysplasia
- Cyclic neutropenia
- PFAPA (periodic fever, aphthae, pharyngitis, adenitis)
- MAGIC (mouth and genital ulcers with inflamed cartilage)

Clinical and Laboratory Workup

-
-
-
-
- capacity
- Folate Vitamin B₁₂, Vitamin B₁, Vitamin B₂, Vitamin B₆
- Weekly CBC x 5
- Serum anti endomysium antibody and transglutaminase assay (positive in celiac disease)
- Bone marrow biopsy

Histology of RAU

- Biopsy and culture to exclude infection
- Biopsy shows a non-specific neutrophilic vascular reaction
- Leukocytoclastic vasculitis
- IF is negative

**Recurrent Aphthous Ulcers
Treatment**

- Begin with topical steroids
- Ointments or gels for lesions in anterior 1/3 of the mouth
- Elixirs for ulcers on pharynx or soft palate
- Eliminate sodium lauryl sulfate (detergent used as emulsifier & surface cleanser) in tooth pastes and mouth rinses
 - Biotin original plain white
 - Rembrandt all natural SLS free
 - Oragel SLS free

**Recurrent Aphthous Ulcers
Topical Therapy**

- Triamcinolone acetonide 0.1% gel or ointment
Fluocinonide 0.05% gel or ointment
Clobetasol 0.05% gel or ointment
- 30 – 60 gm
- Apply a thin film to affected areas QID and leave in place for 30 minutes
- NPO 30-60 minutes

**Dexamethasone
(Decadron, Dexasone)**

- Dexamethasone elixir 0.5 mg/5 mL (high potency, substituted, fluorinated)
- 5 ml QID
- Swish for 5-10 min and expectorate 5 mL QID NPO for 30 min after each dose
- Contact time important for maximizing efficacy; expectorate to limit systemic adverse effects - Cushing syndrome, reversible HPA-axis suppression, hyperglycemia, and glycosuria

Antibiotic Rinses

- Doxycycline 100 mg capsule dissolved in 100 mL H₂O 15 mL Rinse mouth x 5-10 minutes QID
Pregnancy D Avoid < 8 years; Photosensitivity;
Reduce dose in renal impairment; Fanconi-like syndrome may occur with outdated 'ycline
- Chlorhexidine gluconate oral rinse, 0.12% (PerioGard, Peridex) 15 mL swish/spit BID Staining & calculus;
Apply with a Q-tip 11.6% alcohol, saccharin, and mint
- Reduces the severity and pain of ulceration; Not frequency

Coating Agents

- Zilactin-B (OTC) Apply to affected area QID
Contains adhesive gel, 10% benzocaine; tannic, boric and salicylic acids; benzyl alcohol
Zila Pharmaceuticals: (800) 922-7887
- Sucralfate Suspension (an aluminum salt of sucrose octa sulfate) 5 ml QID
Statistically significant improvement in frequency, healing time, and pain scores associated with oral ulcers of Behçet's disease

Topical Medications Anesthetic and Antiinflammatory

- Xylocaine 2% viscous (Lidocaine)
 - TID-QID
 - Risk of aspiration due to loss gag reflex
- Amlexanox (Aphthasol) oral paste 5%
 - 5 gm
 - Apply ¼ inch strip to mucosal ulcers QID (after meals and HS)
 - Wash hands immediately after applying paste. Burning, transient pain, contact mucositis
 - Reassess if not improved after 10 days

Topical Therapy

- Orabase (OTC)
- 7.5 g
- Apply to affected area BID
- Orabase is often used as a base for topical steroid creams due to its adherence to mucosal surface
- Contains gelatin, pectin, and sodium carboxymethylcellulose in polyethylene and mineral oil gel
- Gritty, odorless and tasteless, nonirritating
- Harmless if swallowed

Recurrent Aphthous Stomatitis Systemic Medications

- Prednisone 5 mg, 10 mg, 20 mg tablets
 - Dependent on diagnosis and clinical judgment
 - 1 mg per kg q am and taper to 5 mg QOD as needed
 - Time needed to treat will vary on underlying disease
 - weeks (aphthae)
 - months (pemphigus vulgaris / cicatricial pemphigoid)
- Pentoxifylline (Trental) 400 mg tablets
 - 90 tablets (Therapeutic trial > 30 days)
 - TID with food

Recurrent Aphthous Ulcers Systemic Medications

- Colchicine 0.6 mg tablet
 - 30 tabs (with one refill)
 - 1 tablet po BID to TID
 - Start at BID to minimize GI effects
 - Use in combination with topical and systemic corticosteroids.
 - Adverse effects:* diarrhea, neutropenia, male infertility
 - Contraindications:* renal, GI, cardiac or hematologic disorders

Recurrent Aphthous Ulcers Systemic Medications

- Thalidomide (Thalomid) 50, 100, 200 mg
 - 200 mg PO QD for 4 weeks (gradually increase dose due to sedation)
 - Restricted access in US
 - 1-888-423-5436 or www.thalomid.com
 - FDA approved for HIV associated RAU
 - Off-label use has documented a 55% cure rate compared to 7% in control patients
 - Somnolence, dizziness, constipation, peripheral neuropathy
 - Teratogenic Pregnancy X
- NEJM 1997; 336:1487-1493

Candidiasis

- Common yeast organism
- Normal oral flora
- *Candidiasis* encompasses mucosal and cutaneous conditions
- Oral manifestations: acute or chronic with variable degrees of severity
- Tenderness, burning, dry mouth, thick tongue, pain with swallowing (dysphagia)

Candidiasis Clinical Presentations

- Thrush / pseudomembranous
 - removable, white, plaques on a red base
 - buccal mucosa, tongue, esophagus
- Erythematous candidiasis
 - red thinned mucosa
 - dorsal tongue and palate
- Angular cheilitis
 - Ill fitting dentures --> overclosure --> saliva pooling in the corners of the mouth
- Hypertrophic / median rhomboid glossitis
 - Red and white patches or plaques on dorsal tongue

Pseudomembranous Candidiasis

- White plaques on buccal mucosa, tongue, palate
- Sore mouth
- Swollen lips
- Dry cottony mouth
- Removable

Candidiasis

- Predisposing factors
 - Newborns (sterile gut at birth)
 - Steroids: topical and systemic
 - Antibiotics
 - Diabetes
 - Immunosuppression: chemotherapy , AIDS
 - Dentures, partials, appliances

Median Rhomboid Glossitis

- Shiny oval or diamond shaped elevation, midline, directly in front of the circumvallate papillae.
- *Candida* species may be present
- Histology: Chronic inflammation with fibrosis with occasional hyphae in areas of parakeratosis
- Systemic antifungals helpful

Candidiasis
Topical Treatment*

- 1 po 5x/d x 2 wks
Pitfalls: High sugar content → high caries potential xerostomia; diabetic patients)
- Clotrimazole vaginal tab
dissolve 1 po qhs x 5-7 d
Pitfalls: taste is “disgusting”
- Nystatin oral suspension 600,000 U
swab mouth QID

*

Candidiasis
Systemic Treatment

- [®] (fluconazole)
ii po stat then i po qd x 5 d; then i po M,TH
- Pitfalls: Dentures and appliances
Soak in dilute bleach solutions
(1 t/C water) BID
- Renal dosing (CrCl<50: give usual loading
dose x1, then decrease by 50 %)

Clinical Case

- Elderly man
- Prior hx of lung cancer
- Presented for routine oral examination
- Oral examination reported to be “WNL”

Clinical Case

- Clinical Case: 34 Year Old Male with Fever, A Rash and Acute Enanthem
- D/Dx: Drug reaction, variety of viral illnesses, secondary syphilis, meningococemia, rickettsial diseases
- Dx → Acute HIV Seroconversion

Acute HIV Seroconversion

- distress
- Generalized adenitis, night sweats
- Exanthem: within 1-2 days on trunk and face, morbilliform eruption or hemorrhagic macules
- Enanthem: erythema, ulcerations & candida
- Leukopenia, CSF lymphocytic pleocytosis, transient thrombopenia and lymphopenia

Diagnosis of Acute HIV Seroconversion

-
-
- transmission or large viral inoculum)
- Documentation of laboratory seroconversion
- Circulating p24 antibodies
- D/Dx: Drug reaction, other viral illness, secondary syphilis, meningococemia, rickettsial diseases

Summary

- Oral exam is an integral aspect of the clinical and dermatologic evaluation
- Benign and malignant lesions easily visualized
- Findings reflect wide differential including local and systemic processes
- Now you have knowledge and tools to diagnose and treat these complex patients
- Success is possible and very much achievable

Contact Information

- Wintermeyer Mirowski, DMD, MD
- gmirowsk@iupui.edu

- Private practice
1910 N Arlington St
Indianapolis, IN 46032

- 317-359-5357 Tel
- 317-359-5358 Fax

Treatment Of Mucosal Disorders

- Goal: Maximize success and limit obstacles
- Schedule adequate time
- Remain non-judgmental and supportive
- Obtain a definitive diagnosis
- Avoid telephone diagnoses
 - Reexamine patient
 - Obtain cultures
 - Biopsy
- Treat one condition at a time

Approach to the Patient
What to do after the examination

- Communicate effectively with the patient
- Position the patient and yourself comfortably throughout the examination
- Have patient sit up, return glasses, clothing and any oral appliances before discussing your findings and treatment options

Approach to the Patient

- Identify potential etiologies and exacerbating factors
 - Medical history and review of systems
 - Identify extent of involvement and impact
 - Treat and eliminate confounding conditions
 - Identify potential local or systemic irritants
 - Medications both prescription and OTC
 -
 -
 -
